

# PEROXISOMAL DISORDER FACTS

## NEW FACTS COLLECTED FROM THE EL-PFDD MEETING FOR PEROXISOMAL DISORDERS



01

Peroxisomal disorder patients are often dealing with eleven to twelve symptoms at one time



02

Peroxisomal disorder patients, on average, are administered thirteen to fourteen treatments at one time to manage symptoms



03

Hypotonia, vision loss, hearing loss and intellectual disability are the most frequently reported symptoms in peroxisomal disorders



04

Vision loss is often ranked as the most impactful symptom in peroxisomal disorders



05

Other symptoms that rate high in impact include mobility and intellectual disability



06

The highest ranked (and rated) impact in peroxisomal disorders is their effect on independent activities of daily living

PEROXISOMAL DISORDERS INCLUDED: ZELLWEGER SPECTRUM DISORDER (PREVIOUSLY KNOWN AS ZS, IRD, NALD, AND HEIMLER SYNDROME), D- BIFUNCTIONAL PROTEIN DEFICIENCY, AMACR DEFICIENCY, AND ACDB5 DEFICIENCY.

Mousumi Bose, PhD  
Pamela Saikali, MS  
Montclair State University



GLOBAL  
FOUNDATION  
•for•  
PEROXISOMAL  
DISORDERS