

PEROXISOMAL DISORDER FACTS

TREATMENT FACTS COLLECTED FROM THE EL-PFDD MEETING FOR PEROXISOMAL DISORDERS



01

PD patients, on average, are administered 13-14 treatments at one time to manage symptoms



02

PD treatments encompass at least 5 distinct treatment categories, including: medications, physical devices, therapies/accommodations, supplementations and medical interventions



03

Mobility and sensory impairment are the most frequently treated symptoms in PD



04

Several symptom categories (like GI symptoms) utilize many types of treatments



05

Most treatments for PD have limited effectiveness or lose effectiveness over time



06

Some treatments for PD are inaccessible due to impairment of other symptoms



07

Most treatments for PD have considerable side effects, often exacerbating other symptoms



08

There is a need for new treatments in PD

PEROXISOMAL DISORDERS INCLUDED: ZELLWEGER SPECTRUM DISORDER (PREVIOUSLY KNOWN AS ZS, IRD, NALD, AND HEIMLER SYNDROME), D- BIFUNCTIONAL PROTEIN DEFICIENCY, AMACR DEFICIENCY, AND ACDB5 DEFICIENCY.

Mousumi Bose, PhD
Nicole Fontes, MS
Pamela Saikali, MS
Montclair State University



GLOBAL
FOUNDATION
for
PEROXISOMAL
DISORDERS